

MEAN AND VARIANCE ESTIMATION BY KRIGING

T. SUSLO

ABSTRACT. The aim of the paper is to derive the numerical least-squares estimator for mean and variance of random variable. In order to do so the following questions have to be answered: (i) what is the statistical model for the estimation procedure? (ii) what are the properties of the estimator, like optimality (in which class) or asymptotic properties? (iii) how does the estimator work in practice, how compared to competing estimators?

1. INTRODUCTION

The kriging model based on the stationary random process $\mathcal{V} = \{V_j; \mathbb{N}_1 \ni j \supset i = 1, \dots, n\}$ with the estimation statistics $\hat{V}_j = \omega_j^i V_i$ with an unknown constant mean m and variance σ^2 and some correlation function ρ for the asymptotic solution

$$\lim_{j \rightarrow \infty} E\{[(V_j - m) - (\hat{V}_j - m)]^2\} = \sigma^2$$

has the well-known (co-ordinate independent) least-squares disjunction. The aim of the paper is to find (on computer) a co-ordinate dependent disjunction of kriging model for non-asymptotic solution

$$E\{[(V_j - m) - (\hat{V}_j - m)]^2\} = \sigma^2 .$$

2. CORRELATION FUNCTION ESTIMATOR

Following

$$\begin{aligned} \gamma(h) = \frac{1}{2}E\{(V_j - V_{j+h})^2\} &= \frac{1}{2}E\{V_j^2 - 2V_jV_{j+h} + V_{j+h}^2\} \\ &= E\{V_j^2\} - E\{V_jV_{j+h}\} \\ &= E\{V_j^2\} - E^2\{V_j\} - (E\{V_jV_{j+h}\} - E^2\{V_j\}) \\ &= \sigma^2 - C(h) \geq 0 \end{aligned}$$

we get

$$0 \leq C(h)/C(0) = C(h)/\sigma^2 = 1 - \gamma(h)/\sigma^2 = |\rho(h)| .$$

Since the correlation function estimator must be non-increasing

$$|\hat{\rho}_n(h)| = 1 - \hat{\gamma}_n(h)/\hat{\sigma}^2$$

then only non-decreasing outcomes of the experimental semi-variogram

$$\hat{\gamma}_n(h) = \frac{1}{2} \frac{1}{(n-h)} \sum_{j=1}^{n-h} (v_j - v_{j+h})^2 \quad h = 0, \dots, n-1$$

should be taking into consideration for $h \leq d$

$$\hat{\sigma}^2 = \hat{\gamma}_n(d) \geq \hat{\gamma}_n(d-1) \geq \hat{\gamma}_n(d-2) \geq \dots \geq \hat{\gamma}_n(1) > \hat{\gamma}_n(0) = 0.$$

Since

$$C(h) = E\{V_j V_{j+h}\} - E\{V_j\}E\{V_{j+h}\}$$

then

$$\hat{C}_n(h) = \frac{1}{n-h} \sum_{j=1}^{n-h} v_j v_{j+h} - \frac{1}{(n-h)^2} \sum_{j=1}^{n-h} v_j \sum_{j=1}^{n-h} v_{j+h}$$

and the correlation function estimator is also bounded by

$$|\hat{\rho}_n(h)| = \hat{C}_n(h)/\hat{C}_n(0) \quad h = 0, \dots, d.$$

3. CENTRAL LIMIT THEOREM

Let us consider a set of frozen in time t stationary random processes

$$\mathcal{V}_t = \{V_j; \mathbb{N}_1 \ni j \supset i = 1, \dots, n\}_t; \quad t \in \mathbb{N}_{n+1}$$

since from the Central Limit Theorem holds

$$\begin{aligned} U_{t=n+1} &= \frac{\left[\frac{1}{n} \sum V_j\right]_{t=n+1} - E\{V_j\}_{t=n+1}}{\sqrt{E\{V_j^2\}_{t=n+1} - E^2\{V_j\}_{t=n+1}}} \sqrt{n} \Rightarrow N(0; 1) \\ &\vdots \\ U_{t=n+k} &= \frac{\left[\frac{1}{n} \sum V_j\right]_{t=n+k} - E\{V_j\}_{t=n+k}}{\sqrt{E\{V_j^2\}_{t=n+k} - E^2\{V_j\}_{t=n+k}}} \sqrt{n} \Rightarrow N(0; 1) \end{aligned}$$

an independent set of theirs outcomes

$$u_t = \left[\frac{\bar{v}_j - \omega_j^i v_i}{\sqrt{\omega_j^i v_i^2 - (\omega_j^i v_i)^2}} \sqrt{n} \right]_t$$

for

$$t = n+1, \dots, n+k$$

also must follow the Standard Normal distribution.

Example. Let us consider long-lived asymmetric index profile recorded by 359 close quotes of Warsaw's Stock Market Index shown in Fig. 1. Let the values $v_j = v_{115+1}, \dots, v_{359}$ on the latter asymmetric side be supposed to be response averages $[\bar{v}_j]_t$ of frozen in time t stationary random processes based on $v_i = v_1, \dots, v_{115}$

$$\mathcal{V}_t = \{V_j; \mathbb{N}_1 \ni j \supset i = 1, \dots, 115\}_t; \quad t \in \mathbb{N}_{115+1}$$

with the correlation function

$$\rho_t(\Delta_{ij}) = \begin{cases} -1 \cdot t^{-1.0135} [\Delta_{ij}/t]^2, & \text{for } \Delta_{ij} > 0, \\ +1, & \text{for } \Delta_{ij} = 0, \end{cases}$$




FIGURE 1. Long-lived asymmetric index profile, Warsaw's Stock Market Index (WIG) from 11 X 1999 up to 19 III 2001 (359 close quotes) the numerical least-squares estimator of mean at $j \geq n + 1$ (black dots) compared for frozen model to the asymptotic generalized least-squares estimator of mean (grey line). The dashed vertical line represents $j = n = 115$.

with unknown mean m_t and variance σ_t^2 solving on computer the least-squares constraint in one unknown j for every t

$$\left[E\{[(V_j - m) - (\hat{V}_j - m)]^2\} = \sigma^2 \right]_t ,$$

substituting the solution into

$$\hat{u}_t = \left[\frac{v_j - \omega_j^i v_i}{\sqrt{\omega_j^i v_i^2 - (\omega_j^i v_i)^2}} \sqrt{115} \right]_t$$

we get, for every $t = 115 + 1, \dots, 115 + 102$, the Significance Level for computer K-S test

$$\alpha < 0.19 .$$

REFERENCES

- [1] E. H. Isaaks and R. M. Srivastava, *An Introduction to Applied Geostatistics*, New York: Oxford Univ. Press (1989) 279–307.
- [2] T. Suslo, *Modern Statistics by Kriging*, arXiv:cs.NA/0609079.